



GP guide for imaging patients with scrotal/testicular signs and symptoms

Ultrasound is the first-choice investigation for a range of clinical conditions, including:

- Pain
- Swelling
- Masses

Where possible the GP should examine the patient before referring for a scan.

Referral Guidelines

Presenting Complaint	Imaging Guidance
Palpable lump/mass	<p>Any male with a palpable scrotal lump should have an ultrasound scan. Ultrasound is a helpful investigation to distinguish the characteristics and location (extra - or intratesticular) of the lump.</p> <p>An urgent specialist referral will be recommended under the 14 day pathway for any suspicious masses.</p>
Pain or/and swelling	<p>Ultrasound is helpful to distinguish the following: hydrocoele, epididymitis, orchitis, epididymo-orchitis, spermatocele, varicocele and epididymal cyst.</p>

How do I refer a patient?

To refer a patient for an ultrasound complete one of our referral forms available from our website (www.inhealthgroup.com).

References

iRefer. Royal College of Radiologists. 7th edition, 2012.

Right test, right time, right place; a Framework for Primary Care Access to Imaging. Royal College of Radiologists and Royal College of General Practitioners, 2006.

For further information please call **0844 581 0301**
 email info@inhealthgroup.com or visit www.inhealthgroup.com